arXiv:math-ph/0505085v2 2 Jun 2005

# A Note on the $p \to q$ norms of Completely Positive Maps

Koenraad M.R. Audenaert*
*Blackett Laboratory, Imperial College London, Prince Consort Road, London SW7 2BW, United Kingdom and*
*Institute for Mathematical Sciences, Imperial College London, Exhibition Road, London SW7 2BW, United Kingdom*

(Dated: February 7, 2008)

King and Ruskai asked whether the $p \to q$ norm of a completely positive map $\Phi$, acting between Schatten $p$ and $q$ classes of self-adjoint operators,

$$||\Phi||_{p\to q} = \max_{A=A^*} \frac{||\Phi(A)||_q}{||A||_p},$$

is equal to the $p \to q$ norm of that map when acting between Schatten classes of general, not necessarily self-adjoint, operators. The first proof has been given by Watrous. We give an alternative proof of this statement.

The Schatten $q$-norm is the non-commutative generalisation of the $\ell_q$ norms: for an operator $A$ and for $q \geq 1$

$$||A||_q := (\mathrm{Tr}(|A|^q))^{1/q},$$

where the absolute value $|A|$ is defined as

$$|A| = (A^*A)^{1/2}.$$

The maximal output purity of a completely positive linear map $\Phi$, as measured by the Schatten $q$-norm, is defined as

$$\nu_p(\Phi) = \max_{\substack{A>0 \\ \mathrm{Tr}(A)=1}} ||\Phi(A)||_q.$$

This quantity is a measure of "noisiness" of the map [1], considering it as a channel for the quantum state $A$. It expresses the fact that a pure state will acquire at least this amount of mixedness when passing through the channel.

It has been proven in [1] that this quantity is equal to the $1 \to q$ norm of $\Phi$, considering it as a map from the Schatten 1-class of self-adjoint operators to the Schatten $q$-class. That is

$$\nu_p(\Phi) = \max_{A=A^*} \frac{||\Phi(A)||_q}{||A||_1},$$

where the condition of positivity of $A$ is no longer required.

More generally, one can define the $p \to q$ norm as

$$||\Phi||_{p\to q} := \max_{A=A^*} \frac{||\Phi(A)||_q}{||A||_p}.$$

King and Ruskai asked in [3] whether the condition of self-adjointness of $A$ in this definition can also be dropped, that is, whether the following holds:

$$||\Phi||_{p\to q} = \max_{A} \frac{||\Phi(A)||_q}{||A||_p}.$$

They have proved this in [3] for the case $p = q = 2$, and for the case of completely positive trace preserving maps between 2-dimensional Schatten classes with $p = 1$ and $q \geq 2$.

A completely general proof has been given by Watrous [4] for any completely positive map and for all values of $p, q \geq 1$. In this note we present an alternative proof.

*Proof.* Our proof uses the construction [2] of turning a general operator $A$ on a Hilbert space $\mathcal{H}$ into a self-adjoint one by defining an operator on the doubled space $\mathcal{H} \oplus \mathcal{H}$

$$Q := \begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix},$$

which is obviously self-adjoint.

The absolute value of $A$ is defined as $|A| = (A^*A)^{1/2}$, and there exists a unitary $U$ such that $(AA^*)^{1/2} = U|A|U^*$. The absolute value of $Q$ is therefore

$$\begin{aligned} |Q| &= \left[\begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix} \begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix}\right]^{1/2} \\ &= \begin{pmatrix} AA^* & 0 \\ 0 & A^*A \end{pmatrix}^{1/2} \\ &= \begin{pmatrix} U|A|U^* & 0 \\ 0 & |A| \end{pmatrix}. \end{aligned}$$

The doubled space $\mathcal{H} \oplus \mathcal{H}$ is isomorphic to the space $\mathcal{H} \otimes \mathbb{C}^2$. Let $\Phi \otimes \mathbb{1}_2$ be the completely positive map extending $\Phi$ that operates trivially on $\mathbb{C}^2$. Then, using the fact that for a completely positive map $\Phi(A^*) = \Phi(A)^*$,

$$(\Phi \otimes \mathbb{1}_2)(Q) = \begin{pmatrix} 0 & \Phi(A) \\ \Phi(A)^* & 0 \end{pmatrix}.$$

Thus

$$\begin{aligned} &||(\Phi \otimes \mathbb{1}_2)(Q)||_q^q \\ &= \mathrm{Tr}\, |(\Phi \otimes \mathbb{1}_2)(Q)|^q \\ &= \mathrm{Tr} \left| \begin{pmatrix} 0 & \Phi(A) \\ \Phi(A)^* & 0 \end{pmatrix} \right|^q \\ &= \mathrm{Tr} \begin{pmatrix} (\Phi(A)\Phi(A)^*)^{1/2} & 0 \\ 0 & (\Phi(A)^*\Phi(A))^{1/2} \end{pmatrix}^q \\ &= 2\, \mathrm{Tr}\, |\Phi(A)|^p, \end{aligned}$$

where in the last line we have again used unitary equivalence of $XX^*$ and $X^*X$.

---

[*] Electronic address: k.audenaert@imperial.ac.uk

Likewise,

$$\begin{aligned} ||(\Phi\otimes \mathbb{1}_2)(|Q|)||_q^q &= \mathrm{Tr}(\Phi\otimes \mathbb{1}_2)(|Q|)^q \\ &= \mathrm{Tr}\begin{pmatrix} \Phi(U|A|U^*) & 0 \\ 0 & \Phi(|A|) \end{pmatrix}^q \\ &= \mathrm{Tr}\,\Phi(U|A|U^*)^q + \mathrm{Tr}\,\Phi(|A|)^q. \end{aligned}$$

Here we have used complete positivity of $\Phi$, implying positivity of $(\Phi\otimes \mathbb{1}_2)(|Q|)$.

Amosov and Holevo have shown in [1] that, for any completely positive map $\Omega$, and any self-adjoint operator $X$, $||\Omega(X)||_q \le ||\Omega(|X|)||_q$. Applying this to the map $\Phi\otimes\mathbb{1}_2$ and the operator $Q$, and combining with the above results gives

$$2\,\mathrm{Tr}\,|\Phi(A)|^q \le \mathrm{Tr}\,\Phi(U|A|U^*)^q + \mathrm{Tr}\,\Phi(|A|)^q.$$

Let us now impose the requirement $||A||_p = 1$. Thus, obviously, both $|A|$ and $U|A|U^*$ have $p$-norm equal to 1. Then, by definition of the $p\to q$ norm of $\Phi$,

$$\begin{aligned} \mathrm{Tr}\,\Phi(U|A|U^*)^q &\le ||\Phi||_{p\to q}^q \\ \mathrm{Tr}\,\Phi(|A|)^q &\le ||\Phi||_{p\to q}^q, \end{aligned}$$

so that also

$$2\,\mathrm{Tr}\,|\Phi(A)|^q \le 2||\Phi||_{p\to q}^q.$$

For general $A$, $A/||A||_p$ has $p$-norm equal to 1, so that by homogeneity,

$$\frac{||\Phi(A)||_q}{||A||_p} \le ||\Phi||_{p\to q}.$$

Maximising over all operators $A$, which includes the self-adjoint ones, we find

$$\max_A \frac{||\Phi(A)||_q}{||A||_p} = ||\Phi||_{p\to q}.$$

□

[1] G. G. Amosov and A. S. Holevo, “On the multiplicativity conjecture for quantum channels”, Theor. Probab. Appl. 47, 143146 (2002).

[2] R. Bhatia, *Matrix Analysis*, Springer, Heidelberg (1997).

[3] C. King and M. B. Ruskai, “Comments on multiplicativity of maximal $p$-norms when $p = 2$”, in *Quantum Information, Statistics and Probability*, ed. by O. Hirota, World Scientific; see also quant-ph/0401026 (2004).

[4] J. Watrous, “Notes on super-operator norms induced by Schatten norms”, QIC **5**, 58–68 (2005).